

Toughness and Kronecker Product of Graphs

Authors: Dr. Daniel A. Jaume¹

Departamento de Matemáticas
Universidad Nacional de San Luis, Argentina

—
FoCM 2014 Coference
Montevideo, Uruguay

December 11, 2014

¹djaume@unsl.edu.ar

Vulnerability

Given a connected graph G we ask:

- 1 (minimun) size of a vertex cut set $S \subset V(G)$
- 2 number of remaining connected components $k(G - S)$
- 3 size of the largest connected component $m(G - S)$

Vulnerability

Given a connected graph G we ask:

- 1 (minimum) size of a vertex cut set $S \subset V(G)$
- 2 number of remaining connected components $k(G - S)$
- 3 size of the largest connected component $m(G - S)$

Vulnerability

Given a connected graph G we ask:

- 1 (minimum) size of a vertex cut set $S \subset V(G)$
- 2 number of remaining connected components $k(G - S)$
- 3 size of the largest connected component $m(G - S)$

(Vertex) Vulnerability parameters

192? **connectivity**, $\kappa(G)$, deal with 1

1973 Chvátal's **toughness**, $t(G)$, deal with 1 and 2.

1978 Jung's **scattering number**, $sc(G)$, deal with 1 and 2.

1987 Barefoot-Entringer-Swart's **integrity**, $I(G)$, deal with 1 and 3.

1992 Cozzens-Moazzami-Stueckle's **tenacity**, $T(G)$, deal with 1, 2 and 3.

2004 Li-Zhang-Li's **rupture degree**, $r(G)$, deal with 1, 2 and 3.

(Vertex) Vulnerability parameters

192? **connectivity**, $\kappa(G)$, deal with 1

1973 Chvátal's **toughness**, $t(G)$, deal with 1 and 2.

1978 Jung's **scattering number**, $sc(G)$, deal with 1 and 2.

1987 Barefoot-Entringer-Swart's **integrity**, $I(G)$, deal with 1 and 3.

1992 Cozzens-Moazzami-Stueckle's **tenacity**, $T(G)$, deal with 1, 2 and 3.

2004 Li-Zhang-Li's **rupture degree**, $r(G)$, deal with 1, 2 and 3.

(Vertex) Vulnerability parameters

192? **connectivity**, $\kappa(G)$, deal with 1

1973 Chvátal's **toughness**, $t(G)$, deal with 1 and 2.

1978 Jung's **scattering number**, $sc(G)$, deal with 1 and 2.

1987 Barefoot-Entringer-Swart's **integrity**, $I(G)$, deal with 1 and 3.

1992 Cozzens-Moazzami-Stueckle's **tenacity**, $T(G)$, deal with 1, 2 and 3.

2004 Li-Zhang-Li's **rupture degree**, $r(G)$, deal with 1, 2 and 3.

(Vertex) Vulnerability parameters

192? **connectivity**, $\kappa(G)$, deal with 1

1973 Chvátal's **toughness**, $t(G)$, deal with 1 and 2.

1978 Jung's **scattering number**, $sc(G)$, deal with 1 and 2.

1987 Barefoot-Entringer-Swart's **integrity**, $I(G)$, deal with 1 and 3.

1992 Cozzens-Moazzami-Stueckle's **tenacity**, $T(G)$, deal with 1, 2 and 3.

2004 Li-Zhang-Li's **rupture degree**, $r(G)$, deal with 1, 2 and 3.

(Vertex) Vulnerability parameters

192? **connectivity**, $\kappa(G)$, deal with 1

1973 Chvátal's **toughness**, $t(G)$, deal with 1 and 2.

1978 Jung's **scattering number**, $sc(G)$, deal with 1 and 2.

1987 Barefoot-Entringer-Swart's **integrity**, $I(G)$, deal with 1 and 3.

1992 Cozzens-Moazzami-Stueckle's **tenacity**, $T(G)$, deal with 1, 2 and 3.

2004 Li-Zhang-Li's **rupture degree**, $r(G)$, deal with 1, 2 and 3.

(Vertex) Vulnerability parameters

- 192? **connectivity**, $\kappa(G)$, deal with 1
- 1973 Chvátal's **toughness**, $t(G)$, deal with 1 and 2.
- 1978 Jung's **scattering number**, $sc(G)$, deal with 1 and 2.
- 1987 Barefoot-Entringer-Swart's **integrity**, $I(G)$, deal with 1 and 3.
- 1992 Cozzens-Moazzami-Stueckle's **tenacity**, $T(G)$, deal with 1, 2 and 3.
- 2004 Li-Zhang-Li's **rupture degree**, $r(G)$, deal with 1, 2 and 3.

Toughness

Definition

Given a connected non-complete graph G , the **toughness** of G is:

$$t(G) := \min \frac{|S|}{k(G - S)}$$

where the minimum is taken over all the vertex-cut sets $S \subset V(G)$.
By definition $t(K_n) := +\infty$

Example

- $t(C_n) = 1$, $n \geq 3$.
- $t(K_{k,n-k}) = \frac{k}{n-k}$, where $1 \leq k \leq \frac{n}{2}$.
- $t(\text{Petersen graph}) = \frac{4}{3}$.
- Every Hamiltonian graph have toughness at least 1.

Toughness

Definition

Given a connected non-complete graph G , the **toughness** of G is:

$$t(G) := \min \frac{|S|}{k(G - S)}$$

where the minimum is taken over all the vertex-cut sets $S \subset V(G)$.
By definition $t(K_n) := +\infty$

Example

- $t(C_n) = 1, n \geq 3$.
- $t(K_{k,n-k}) = \frac{k}{n-k}$, where $1 \leq k \leq \frac{n}{2}$.
- $t(\text{Petersen graph}) = \frac{4}{3}$.
- Every Hamiltonian graph have toughness at least 1.

Toughness

Definition

Given a connected non-complete graph G , the **toughness** of G is:

$$t(G) := \min \frac{|S|}{k(G - S)}$$

where the minimum is taken over all the vertex-cut sets $S \subset V(G)$.
By definition $t(K_n) := +\infty$

Example

- $t(C_n) = 1, n \geq 3$.
- $t(K_{k,n-k}) = \frac{k}{n-k}$, where $1 \leq k \leq \frac{n}{2}$.
- $t(\text{Petersen graph}) = \frac{4}{3}$.
- Every Hamiltonian graph have toughness at least 1.

Toughness

Definition

Given a connected non-complete graph G , the **toughness** of G is:

$$t(G) := \min \frac{|S|}{k(G - S)}$$

where the minimum is taken over all the vertex-cut sets $S \subset V(G)$.
By definition $t(K_n) := +\infty$

Example

- $t(C_n) = 1$, $n \geq 3$.
- $t(K_{k,n-k}) = \frac{k}{n-k}$, where $1 \leq k \leq \frac{n}{2}$.
- $t(\text{Petersen graph}) = \frac{4}{3}$.
- Every Hamiltonian graph have toughness at least 1.

Toughness

Definition

Given a connected non-complete graph G , the **toughness** of G is:

$$t(G) := \min \frac{|S|}{k(G - S)}$$

where the minimum is taken over all the vertex-cut sets $S \subset V(G)$.
By definition $t(K_n) := +\infty$

Example

- $t(C_n) = 1$, $n \geq 3$.
- $t(K_{k,n-k}) = \frac{k}{n-k}$, where $1 \leq k \leq \frac{n}{2}$.
- $t(\text{Petersen graph}) = \frac{4}{3}$.
- Every Hamiltonian graph have toughness at least 1.

Scattering number

Definition

Given a connected non-complete graph G , the **scattering number** of G is:

$$sc(G) := \max \{k(G - S) - |S|\}$$

where the maximum is taken over all the vertex-cut sets $S \subset V(G)$. By definition $sc(K_n) := -\infty$

Example

- $sc(C_n) = 0$, for $n \geq 4$.
- $sc(P_n) = 1$, for $n \geq 3$.
- $sc(K_{m,n}) = n - m$, if $m \leq n$ and $n \geq 2$.

Scattering number

Definition

Given a connected non-complete graph G , the **scattering number** of G is:

$$sc(G) := \max \{k(G - S) - |S|\}$$

where the maximum is taken over all the vertex-cut sets $S \subset V(G)$. By definition $sc(K_n) := -\infty$

Example

- $sc(C_n) = 0$, for $n \geq 4$.
- $sc(P_n) = 1$, for $n \geq 3$.
- $sc(K_{m,n}) = n - m$, if $m \leq n$ and $n \geq 2$.

Scattering number

Definition

Given a connected non-complete graph G , the **scattering number** of G is:

$$sc(G) := \max \{k(G - S) - |S|\}$$

where the maximum is taken over all the vertex-cut sets $S \subset V(G)$. By definition $sc(K_n) := -\infty$

Example

- $sc(C_n) = 0$, for $n \geq 4$.
- $sc(P_n) = 1$, for $n \geq 3$.
- $sc(K_{m,n}) = n - m$, if $m \leq n$ and $n \geq 2$.

Scattering number

Definition

Given a connected non-complete graph G , the **scattering number** of G is:

$$sc(G) := \max \{k(G - S) - |S|\}$$

where the maximum is taken over all the vertex-cut sets $S \subset V(G)$. By definition $sc(K_n) := -\infty$

Example

- $sc(C_n) = 0$, for $n \geq 4$.
- $sc(P_n) = 1$, for $n \geq 3$.
- $sc(K_{m,n}) = n - m$, if $m \leq n$ and $n \geq 2$.

Integrity

Definition

Given a connected non-complete graph G , the **integrity** of G is:

$$I(G) := \min\{|S| + m(G - S)\}$$

where the minimum is taken over all the vertex-cut sets $S \subset V(G)$.
By definition $I(K_n) := n$.

Example

- $I(P_n) = \lceil 2\sqrt{n+1} \rceil - 2$.
- $I(K_{m,n}) = 1 + \min\{m, n\}$.

Integrity

Definition

Given a connected non-complete graph G , the **integrity** of G is:

$$I(G) := \min\{|S| + m(G - S)\}$$

where the minimum is taken over all the vertex-cut sets $S \subset V(G)$.
By definition $I(K_n) := n$.

Example

- $I(P_n) = \lceil 2\sqrt{n+1} \rceil - 2$.
- $I(K_{m,n}) = 1 + \min\{m, n\}$.

Integrity

Definition

Given a connected non-complete graph G , the **integrity** of G is:

$$I(G) := \min\{|S| + m(G - S)\}$$

where the minimum is taken over all the vertex-cut sets $S \subset V(G)$.
By definition $I(K_n) := n$.

Example

- $I(P_n) = \lceil 2\sqrt{n+1} \rceil - 2$.
- $I(K_{m,n}) = 1 + \min\{m, n\}$.

Tenacity

Definition

Given a connected non-complete graph G , the **tenacity** of G is

$$T(G) := \min \left\{ \frac{|S| + m(G - S)}{k(G - S)} \right\}$$

where the minimum is taken over all the vertex-cut sets $S \subset V(G)$.
By definition $T(K_n) := n$.

Example

- $T(P_n) = 1$, if n is odd, and $T(P_n) = \frac{n+2}{n}$, if n is even.
- $T(C_n) = \frac{n+3}{n-1}$, if n is odd, and $T(P_n) = \frac{n+2}{n}$, if n is even.
- $T(K_{k,n-k}) = \frac{k+1}{n-k}$, for $1 \leq k \leq \frac{n}{2}$

Tenacity

Definition

Given a connected non-complete graph G , the **tenacity** of G is

$$T(G) := \min \left\{ \frac{|S| + m(G - S)}{k(G - S)} \right\}$$

where the minimum is taken over all the vertex-cut sets $S \subset V(G)$.
By definition $T(K_n) := n$.

Example

- $T(P_n) = 1$, if n is odd, and $T(P_n) = \frac{n+2}{n}$, if n is even.
- $T(C_n) = \frac{n+3}{n-1}$, if n is odd, and $T(P_n) = \frac{n+2}{n}$, if n is even.
- $T(K_{k,n-k}) = \frac{k+1}{n-k}$, for $1 \leq k \leq \frac{n}{2}$

Tenacity

Definition

Given a connected non-complete graph G , the **tenacity** of G is

$$T(G) := \min \left\{ \frac{|S| + m(G - S)}{k(G - S)} \right\}$$

where the minimum is taken over all the vertex-cut sets $S \subset V(G)$.
By definition $T(K_n) := n$.

Example

- $T(P_n) = 1$, if n is odd, and $T(P_n) = \frac{n+2}{n}$, if n is even.
- $T(C_n) = \frac{n+3}{n-1}$, if n is odd, and $T(P_n) = \frac{n+2}{n}$, if n is even.
- $T(K_{k,n-k}) = \frac{k+1}{n-k}$, for $1 \leq k \leq \frac{n}{2}$

Tenacity

Definition

Given a connected non-complete graph G , the **tenacity** of G is

$$T(G) := \min \left\{ \frac{|S| + m(G - S)}{k(G - S)} \right\}$$

where the minimum is taken over all the vertex-cut sets $S \subset V(G)$.
By definition $T(K_n) := n$.

Example

- $T(P_n) = 1$, if n is odd, and $T(P_n) = \frac{n+2}{n}$, if n is even.
- $T(C_n) = \frac{n+3}{n-1}$, if n is odd, and $T(P_n) = \frac{n+2}{n}$, if n is even.
- $T(K_{k,n-k}) = \frac{k+1}{n-k}$, for $1 \leq k \leq \frac{n}{2}$

Rupture degree

Definition

Given a connected non-complete graph G , the **rupture degree** of G is

$$r(G) := \max \{k(G - S) - |S| - m(G - S)\}$$

where the maximum is taken over all the vertex-cut sets $S \subset V(G)$.
By definition $r(K_n) := 1 - n$.

Example

- $r(P_n) = 0$, if n is odd, and $r(P_n) = -1$, if n is even.
- $r(C_n) = -2$, if n is odd, and $r(C_n) = -1$, if n is even.
- $r(K_{k,n-k}) = n - 2k - 1$, for $1 \leq k \leq \frac{n}{2}$

Rupture degree

Definition

Given a connected non-complete graph G , the **rupture degree** of G is

$$r(G) := \max \{k(G - S) - |S| - m(G - S)\}$$

where the maximum is taken over all the vertex-cut sets $S \subset V(G)$.
By definition $r(K_n) := 1 - n$.

Example

- $r(P_n) = 0$, if n is odd, and $r(P_n) = -1$, if n is even.
- $r(C_n) = -2$, if n is odd, and $r(C_n) = -1$, if n is even.
- $r(K_{k,n-k}) = n - 2k - 1$, for $1 \leq k \leq \frac{n}{2}$

Rupture degree

Definition

Given a connected non-complete graph G , the **rupture degree** of G is

$$r(G) := \max \{k(G - S) - |S| - m(G - S)\}$$

where the maximum is taken over all the vertex-cut sets $S \subset V(G)$.
By definition $r(K_n) := 1 - n$.

Example

- $r(P_n) = 0$, if n is odd, and $r(P_n) = -1$, if n is even.
- $r(C_n) = -2$, if n is odd, and $r(C_n) = -1$, if n is even.
- $r(K_{k,n-k}) = n - 2k - 1$, for $1 \leq k \leq \frac{n}{2}$

Rupture degree

Definition

Given a connected non-complete graph G , the **rupture degree** of G is

$$r(G) := \max \{k(G - S) - |S| - m(G - S)\}$$

where the maximum is taken over all the vertex-cut sets $S \subset V(G)$.
By definition $r(K_n) := 1 - n$.

Example

- $r(P_n) = 0$, if n is odd, and $r(P_n) = -1$, if n is even.
- $r(C_n) = -2$, if n is odd, and $r(C_n) = -1$, if n is even.
- $r(K_{k,n-k}) = n - 2k - 1$, for $1 \leq k \leq \frac{n}{2}$

Kronecker Product

Definition

Let G, H be two graphs. The **Kronecker Product** of both is the graph (Nešetřil notation)

$$G \times H$$

with vertex set

$$V(G \times H) = V(G) \times V(H)$$

(Cartesian product of sets) and edge set

$$E(G \times H) = \{ \{(a, b), (a', b')\} \mid \{a, a'\} \in E(G), \{b, b'\} \in E(H) \}$$

Results of Mamut-Vumar 2007, IPL

Theorem

Let n y m be integer with $n \geq m \geq 2$ and $n \geq 3$. Then:

- ① $t(K_m \times K_n) = m - 1$
- ② $sc(K_m \times K_n) = \begin{cases} 2 - (m - 1)(n - 1), & \text{if } m = n \\ 2n - mn, & \text{otherwise} \end{cases}$
- ③ $I(K_m \otimes K_n) = mn - n + 1$
- ④ $T(K_m \otimes K_n) = m + \frac{1}{n} - 1$
- ⑤ $r(K_m \otimes K_n) = ???$

Results of Mamut-Vumar 2007, IPL

Theorem

Let n y m be integer with $n \geq m \geq 2$ and $n \geq 3$. Then:

- ① $t(K_m \times K_n) = m - 1$
- ② $sc(K_m \times K_n) = \begin{cases} 2 - (m - 1)(n - 1), & \text{if } m = n \\ 2n - mn, & \text{otherwise} \end{cases}$
- ③ $I(K_m \otimes K_n) = mn - n + 1$
- ④ $T(K_m \otimes K_n) = m + \frac{1}{n} - 1$
- ⑤ $r(K_m \otimes K_n) = ???$

Results of Mamut-Vumar 2007, IPL

Theorem

Let n y m be integer with $n \geq m \geq 2$ and $n \geq 3$. Then:

- ① $t(K_m \times K_n) = m - 1$
- ② $sc(K_m \times K_n) = \begin{cases} 2 - (m - 1)(n - 1), & \text{if } m = n \\ 2n - mn, & \text{otherwise} \end{cases}$
- ③ $I(K_m \otimes K_n) = mn - n + 1$
- ④ $T(K_m \otimes K_n) = m + \frac{1}{n} - 1$
- ⑤ $r(K_m \otimes K_n) = ???$

Results of Mamut-Vumar 2007, IPL

Theorem

Let n y m be integer with $n \geq m \geq 2$ and $n \geq 3$. Then:

- 1 $t(K_m \times K_n) = m - 1$
- 2 $sc(K_m \times K_n) = \begin{cases} 2 - (m - 1)(n - 1), & \text{if } m = n \\ 2n - mn, & \text{otherwise} \end{cases}$
- 3 $I(K_m \otimes K_n) = mn - n + 1$
- 4 $T(K_m \otimes K_n) = m + \frac{1}{n} - 1$
- 5 $r(K_m \otimes K_n) = ???$

Results of Mamut-Vumar 2007, IPL

Theorem

Let n y m be integer with $n \geq m \geq 2$ and $n \geq 3$. Then:

- ① $t(K_m \times K_n) = m - 1$
- ② $sc(K_m \times K_n) = \begin{cases} 2 - (m - 1)(n - 1), & \text{if } m = n \\ 2n - mn, & \text{otherwise} \end{cases}$
- ③ $I(K_m \otimes K_n) = mn - n + 1$
- ④ $T(K_m \otimes K_n) = m + \frac{1}{n} - 1$
- ⑤ $r(K_m \otimes K_n) = ???$

Mamut-Vumar-Problem

Problem (Mamut y Vumar (2007))

Determine good bounds for vulnerability parameters of $G \times H$, with G and H arbitrary connected graphs.

Our little result

Theorem

Let G be a connected non-complete graph such that $t(G) \geq \frac{n}{2}$, with $n \geq 3$. Then

$$t(G \times K_n) = n - 1$$

Unitary Cayley Graph

Definition


Let n be a positive integer greater than 1. The Unitary Cayley Graph X_n is defined as the graph with vertex set

$$V(X_n) = \{0, 1, \dots, n-1\}$$


and edge set

$$E(X_n) = \{(a, b) \mid a, b \in \mathbb{Z}_n, \quad \gcd(a - b, n) = 1\}$$


Example: X_{10}




$$U_{10} = \{1, 3, 7, 9\}$$

Example: X_{10} 


$$U_{10} = \{1, 3, 7, 9\}$$

Example: X_{10} 

$$U_{10} = \{1, 3, 7, 9\}$$

Example: X_{10} 

$$U_{10} = \{1, 3, 7, 9\}$$

Example: X_{10} 

$$U_{10} = \{1, 3, 7, 9\}$$

Results by Klotz and Sander 2007, EJC

Some properties of unitary Cayley Graphs were determined by Klotz and Sander [4].

Here, $p_1(n)$ will denote the smallest prime number p such that $p|n$.

Theorem

Let X_n be a Unitary Cayley Graph of order n then

- ① *the independence number $\text{ind}(X_n) = \frac{n}{p_1(n)}$.*
- ② *the vertex connectivity $\kappa(X_n) = \varphi(n)$.*

Results by Klotz and Sander 2007, EJC

Some properties of unitary Cayley Graphs were determined by Klotz and Sander [4].

Here, $p_1(n)$ will denote the smallest prime number p such that $p|n$.

Theorem

Let X_n be a Unitary Cayley Graph of order n then

- ① *the independence number $\text{ind}(X_n) = \frac{n}{p_1(n)}$.*
- ② *the vertex connectivity $\kappa(X_n) = \varphi(n)$.*

Results by Klotz and Sander 2007, EJC

Some properties of unitary Cayley Graphs were determined by Klotz and Sander [4].

Here, $p_1(n)$ will denote the smallest prime number p such that $p|n$.

Theorem

Let X_n be a Unitary Cayley Graph of order n then

- 1 *the independence number $\text{ind}(X_n) = \frac{n}{p_1(n)}$.*
- 2 *the vertex connectivity $\kappa(X_n) = \varphi(n)$.*

Results by Klotz and Sander 2007, EJC

Some properties of unitary Cayley Graphs were determined by Klotz and Sander [4].

Here, $p_1(n)$ will denote the smallest prime number p such that $p|n$.

Theorem

Let X_n be a Unitary Cayley Graph of order n then

- 1 *the independence number $\text{ind}(X_n) = \frac{n}{p_1(n)}$.*
- 2 *the vertex connectivity $\kappa(X_n) = \varphi(n)$.*

Results by Klotz and Sander 2007, EJC

Some properties of unitary Cayley Graphs were determined by Klotz and Sander [4].

Here, $p_1(n)$ will denote the smallest prime number p such that $p|n$.

Theorem

Let X_n be a Unitary Cayley Graph of order n then

- 1 *the independence number $\text{ind}(X_n) = \frac{n}{p_1(n)}$.*
- 2 *the vertex connectivity $\kappa(X_n) = \varphi(n)$.*

One result of Ramaswamy and Veena 2009, EJC

Theorem

Given $n \in \mathbb{Z}^+$, with prime factorization $= p_1^{\alpha_1} \dots p_k^{\alpha_k}$, where $p_1 < \dots < p_k$. Then

$$X_n \cong X_{p_1^{\alpha_1}} \times \dots \times X_{p_k^{\alpha_k}}$$

radical bound

Given $n \in \mathbb{Z}^+$, the **radical** of n , denoted $\text{rad}(n)$, is the greatest square-free divisor of n .

Theorem

Let X_n be an Unitary Cayley Graph of order n . Then
 $t(X_n) \geq t(X_{\text{rad}(n)})$

Proof.

- Take a t -vertex-cut-set $S \subset V(X_n)$.
- $Z := \{i \in \mathbb{Z}_{\text{rad}(n)} \mid \text{if } g \equiv i \pmod{\text{rad}(n)}, \text{ then } g \in S\}$
 - $|S| \geq \frac{n}{\text{rad}(n)} |Z|$.
 - $k(X_n - S) \leq k(X_{\text{rad}(n)} - Z)$
-

$$t(X_n) = \frac{|S|}{k(X_n - S)} \geq \frac{\frac{n}{\text{rad}(n)} |Z|}{\frac{n}{\text{rad}(n)} k(X_{\text{rad}(n)} - Z)} \geq t(X_{\text{rad}(n)})$$

radical bound

Given $n \in \mathbb{Z}^+$, the **radical** of n , denoted $\text{rad}(n)$, is the greatest square-free divisor of n .

Theorem

Let X_n be an Unitary Cayley Graph of order n . Then
 $t(X_n) \geq t(X_{\text{rad}(n)})$

Proof.

- Take a t -vertex-cut-set $S \subset V(X_n)$.
- $Z := \{i \in \mathbb{Z}_{\text{rad}(n)} \mid \text{if } g \equiv i \pmod{\text{rad}(n)}, \text{ then } g \in S\}$
 - $|S| \geq \frac{n}{\text{rad}(n)} |Z|$.
 - $k(X_n - S) \leq k(X_{\text{rad}(n)} - Z)$
-

$$t(X_n) = \frac{|S|}{k(X_n - S)} \geq \frac{\frac{n}{\text{rad}(n)} |Z|}{\frac{n}{\text{rad}(n)} k(X_{\text{rad}(n)} - Z)} \geq t(X_{\text{rad}(n)})$$

radical bound

Given $n \in \mathbb{Z}^+$, the **radical** of n , denoted $\text{rad}(n)$, is the greatest square-free divisor of n .

Theorem

Let X_n be an Unitary Cayley Graph of order n . Then
 $t(X_n) \geq t(X_{\text{rad}(n)})$

Proof.

- Take a t -vertex-cut-set $S \subset V(X_n)$.
- $Z := \{i \in \mathbb{Z}_{\text{rad}(n)} \mid \text{if } g \equiv i \pmod{\text{rad}(n)}, \text{ then } g \in S\}$
 - $|S| \geq \frac{n}{\text{rad}(n)} |Z|$.
 - $k(X_n - S) \leq k(X_{\text{rad}(n)} - Z)$
-

$$t(X_n) = \frac{|S|}{k(X_n - S)} \geq \frac{\frac{n}{\text{rad}(n)} |Z|}{\frac{n}{\text{rad}(n)} k(X_{\text{rad}(n)} - Z)} \geq t(X_{\text{rad}(n)})$$

radical bound

Given $n \in \mathbb{Z}^+$, the **radical** of n , denoted $\text{rad}(n)$, is the greatest square-free divisor of n .

Theorem

Let X_n be an Unitary Cayley Graph of order n . Then
 $t(X_n) \geq t(X_{\text{rad}(n)})$

Proof.

- Take a t -vertex-cut-set $S \subset V(X_n)$.
- $Z := \{i \in \mathbb{Z}_{\text{rad}(n)} \mid \text{if } g \equiv i \pmod{\text{rad}(n)}, \text{ then } g \in S\}$
 - $|S| \geq \frac{n}{\text{rad}(n)} |Z|$.
 - $k(X_n - S) \leq k(X_{\text{rad}(n)} - Z)$
-

$$t(X_n) = \frac{|S|}{k(X_n - S)} \geq \frac{\frac{n}{\text{rad}(n)} |Z|}{\frac{n}{\text{rad}(n)} k(X_{\text{rad}(n)} - Z)} \geq t(X_{\text{rad}(n)})$$

radical bound

Given $n \in \mathbb{Z}^+$, the **radical** of n , denoted $\text{rad}(n)$, is the greatest square-free divisor of n .

Theorem

Let X_n be an Unitary Cayley Graph of order n . Then
 $t(X_n) \geq t(X_{\text{rad}(n)})$

Proof.

- Take a t -vertex-cut-set $S \subset V(X_n)$.
- $Z := \{i \in \mathbb{Z}_{\text{rad}(n)} \mid \text{if } g \equiv i \pmod{\text{rad}(n)}, \text{ then } g \in S\}$
 - $|S| \geq \frac{n}{\text{rad}(n)} |Z|$.
 - $k(X_n - S) \leq k(X_{\text{rad}(n)} - Z)$

•

$$t(X_n) = \frac{|S|}{k(X_n - S)} \geq \frac{\frac{n}{\text{rad}(n)} |Z|}{\frac{n}{\text{rad}(n)} k(X_{\text{rad}(n)} - Z)} \geq t(X_{\text{rad}(n)})$$

radical bound

Given $n \in \mathbb{Z}^+$, the **radical** of n , denoted $\text{rad}(n)$, is the greatest square-free divisor of n .

Theorem

Let X_n be an Unitary Cayley Graph of order n . Then
 $t(X_n) \geq t(X_{\text{rad}(n)})$

Proof.

- Take a t -vertex-cut-set $S \subset V(X_n)$.
- $Z := \{i \in \mathbb{Z}_{\text{rad}(n)} \mid \text{if } g \equiv i \pmod{\text{rad}(n)}, \text{ then } g \in S\}$
 - $|S| \geq \frac{n}{\text{rad}(n)} |Z|$.
 - $k(X_n - S) \leq k(X_{\text{rad}(n)} - Z)$
-

$$t(X_n) = \frac{|S|}{k(X_n - S)} \geq \frac{\frac{n}{\text{rad}(n)} |Z|}{\frac{n}{\text{rad}(n)} k(X_{\text{rad}(n)} - Z)} \geq t(X_{\text{rad}(n)})$$

Tools

First a very well known result about toughness

Theorem

For every noncomplete graph G ,

$$\frac{\kappa(G)}{\text{ind}(G)} \leq t(G) \leq \frac{\kappa(G)}{2}$$

Now, a humble result of us:

Theorem

Let $p < q$ be prime numbers and S vertex-cut set with $|S| < pq - q$ then $k(X_{pq} - S) = 2$.

Tools

First a very well known result about toughness

Theorem

For every noncomplete graph G ,

$$\frac{\kappa(G)}{\text{ind}(G)} \leq t(G) \leq \frac{\kappa(G)}{2}$$

Now, a humble result of us:

Theorem

Let $p < q$ be prime numbers and S vertex-cut set with $|S| < pq - q$ then $k(X_{pq} - S) = 2$.

Tools

First a very well known result about toughness

Theorem


For every noncomplete graph G ,


$$\frac{\kappa(G)}{\text{ind}(G)} \leq t(G) \leq \frac{\kappa(G)}{2}$$


Now, a humble result of us:


Theorem


Let $p < q$ be prime numbers and S vertex-cut set with $|S| < pq - q$ then $k(X_{pq} - S) = 2$.











Toughness of UCG

Theorem

Let X_n be an Unitary Cayley Graph of order n . Then
 $t(X_n) = p_1(n) - 1$

Proof.

Assume that n is square-free.

Induction over the number of prime divisor of n

- $n = p_1 p_2$, and $S \subset V(X_n)$ a vertex-cut of X_n
- If $|S| < p_1 p_2 - p_2$, then $k(X_{p_1 p_2} - S) = 2$.
-

$$\frac{|S|}{k(X_{p_1 p_2} - S)} \geq \frac{\kappa(X_{p_1 p_2})}{2} = \frac{(p_1 - 1)(p_2 - 1)}{2} \geq p_1 - 1$$



Toughness of UCG

Theorem

Let X_n be an Unitary Cayley Graph of order n . Then
 $t(X_n) = p_1(n) - 1$

Proof.

Assume that n is square-free.

Induction over the number of prime divisor of n

- $n = p_1 p_2$, and $S \subset V(X_n)$ a vertex-cut of X_n
- If $|S| < p_1 p_2 - p_2$, then $k(X_{p_1 p_2} - S) = 2$.
-

$$\frac{|S|}{k(X_{p_1 p_2} - S)} \geq \frac{\kappa(X_{p_1 p_2})}{2} = \frac{(p_1 - 1)(p_2 - 1)}{2} \geq p_1 - 1$$



Toughness of UCG

Theorem

Let X_n be an Unitary Cayley Graph of order n . Then
 $t(X_n) = p_1(n) - 1$

Proof.

Assume that n is square-free.

Induction over the number of prime divisor of n

- $n = p_1 p_2$, and $S \subset V(X_n)$ a vertex-cut of X_n
- If $|S| < p_1 p_2 - p_2$, then $k(X_{p_1 p_2} - S) = 2$.

•

$$\frac{|S|}{k(X_{p_1 p_2} - S)} \geq \frac{\kappa(X_{p_1 p_2})}{2} = \frac{(p_1 - 1)(p_2 - 1)}{2} \geq p_1 - 1$$



Toughness of UCG

Theorem

Let X_n be an Unitary Cayley Graph of order n . Then
 $t(X_n) = p_1(n) - 1$

Proof.

Assume that n is square-free.

Induction over the number of prime divisor of n

- $n = p_1 p_2$, and $S \subset V(X_n)$ a vertex-cut of X_n
- If $|S| < p_1 p_2 - p_2$, then $k(X_{p_1 p_2} - S) = 2$.
-

$$\frac{|S|}{k(X_{p_1 p_2} - S)} \geq \frac{\kappa(X_{p_1 p_2})}{2} = \frac{(p_1 - 1)(p_2 - 1)}{2} \geq p_1 - 1$$



proof

Proof.

- If $|S| \geq p_1 p_2 - p_2$, then

$$\frac{|S|}{k(X_{p_1 p_2} - S)} \geq \frac{p_1 p_2 - p_2}{\text{ind}(X_{p_1 p_2})} = p_1 - 1$$

- $n = p_1 p_2 \dots p_{k+1}$ con $p_1 < \dots < p_{k+1}$
- $X_n \cong X_{p_1} \times G$, where $G = X_{p_2} \times \dots \times X_{p_{k+1}}$
- But $t(G) \geq p_2 - 1$ by induction hypothesis.
- Then, $t(X_{p_1} \times G) = p_1 - 1$.
- If n is non-square-free, as $t(X_{\text{rad}(n)}) \leq t(X_n) \leq p_1(n) - 1$, we are done!.



proof

Proof.

- If $|S| \geq p_1 p_2 - p_2$, then

$$\frac{|S|}{k(X_{p_1 p_2} - S)} \geq \frac{p_1 p_2 - p_2}{\text{ind}(X_{p_1 p_2})} = p_1 - 1$$

- $n = p_1 p_2 \dots p_{k+1}$ con $p_1 < \dots < p_{k+1}$
- $X_n \cong X_{p_1} \times G$, where $G = X_{p_2} \times \dots \times X_{p_{k+1}}$
- But $t(G) \geq p_2 - 1$ by induction hypothesis.
- Then, $t(X_{p_1} \times G) = p_1 - 1$.
- If n is non-square-free, as $t(X_{\text{rad}(n)}) \leq t(X_n) \leq p_1(n) - 1$, we are done!.



proof

Proof.

- If $|S| \geq p_1 p_2 - p_2$, then

$$\frac{|S|}{k(X_{p_1 p_2} - S)} \geq \frac{p_1 p_2 - p_2}{\text{ind}(X_{p_1 p_2})} = p_1 - 1$$

- $n = p_1 p_2 \dots p_{k+1}$ con $p_1 < \dots < p_{k+1}$
- $X_n \cong X_{p_1} \times G$, where $G = X_{p_2} \times \dots \times X_{p_{k+1}}$
- But $t(G) \geq p_2 - 1$ by induction hypothesis.
- Then, $t(X_{p_1} \times G) = p_1 - 1$.
- If n is non-square-free, as $t(X_{\text{rad}(n)}) \leq t(X_n) \leq p_1(n) - 1$, we are done!.



proof

Proof.

- If $|S| \geq p_1 p_2 - p_2$, then

$$\frac{|S|}{k(X_{p_1 p_2} - S)} \geq \frac{p_1 p_2 - p_2}{\text{ind}(X_{p_1 p_2})} = p_1 - 1$$

- $n = p_1 p_2 \dots p_{k+1}$ con $p_1 < \dots < p_{k+1}$
- $X_n \cong X_{p_1} \times G$, where $G = X_{p_2} \times \dots \times X_{p_{k+1}}$
- But $t(G) \geq p_2 - 1$ by induction hypothesis.
- Then, $t(X_{p_1} \times G) = p_1 - 1$.
- If n is non-square-free, as $t(X_{\text{rad}(n)}) \leq t(X_n) \leq p_1(n) - 1$, we are done!.



proof

Proof.

- If $|S| \geq p_1 p_2 - p_2$, then

$$\frac{|S|}{k(X_{p_1 p_2} - S)} \geq \frac{p_1 p_2 - p_2}{\text{ind}(X_{p_1 p_2})} = p_1 - 1$$

- $n = p_1 p_2 \dots p_{k+1}$ con $p_1 < \dots < p_{k+1}$
- $X_n \cong X_{p_1} \times G$, where $G = X_{p_2} \times \dots \times X_{p_{k+1}}$
- But $t(G) \geq p_2 - 1$ by induction hypothesis.
- Then, $t(X_{p_1} \times G) = p_1 - 1$.
- If n is non-square-free, as $t(X_{\text{rad}(n)}) \leq t(X_n) \leq p_1(n) - 1$, we are done!.



proof

Proof.

- If $|S| \geq p_1 p_2 - p_2$, then

$$\frac{|S|}{k(X_{p_1 p_2} - S)} \geq \frac{p_1 p_2 - p_2}{\text{ind}(X_{p_1 p_2})} = p_1 - 1$$

- $n = p_1 p_2 \dots p_{k+1}$ con $p_1 < \dots < p_{k+1}$
- $X_n \cong X_{p_1} \times G$, where $G = X_{p_2} \times \dots \times X_{p_{k+1}}$
- But $t(G) \geq p_2 - 1$ by induction hypothesis.
- Then, $t(X_{p_1} \times G) = p_1 - 1$.
- If n is non-square-free, as $t(X_{\text{rad}(n)}) \leq t(X_n) \leq p_1(n) - 1$, we are done!.



Tenacity

It is well known that:

Theorem

Let G be a non-complete connected graph. Then

$$t(G) + \frac{1}{\text{ind}(G)} \leq T(G) \leq \frac{|V(G)| - \text{ind}(G) + 1}{\text{ind}(G)}$$

With this we can prove that

Corollary

Let X_n a Unitary Cayley Graph of order n . Then

$$T(X_n) = p_1(n) \left(1 + \frac{1}{n} \right) - 1$$

Tenacity

It is well known that:

Theorem

Let G be a non-complete connected graph. Then

$$t(G) + \frac{1}{\text{ind}(G)} \leq T(G) \leq \frac{|V(G)| - \text{ind}(G) + 1}{\text{ind}(G)}$$

With this we can prove that

Corollary

Let X_n a Unitary Cayley Graph of order n . Then

$$T(X_n) = p_1(n) \left(1 + \frac{1}{n} \right) - 1$$

Integrity

Remember

Definition

Given a non-complete connected graph G , the **integrity** of G , $I(G)$ is defined as:

$$\min\{|S| + m(G - S)\}$$

where the minimum is taken over all the $S \subset V(G)$.

A Known Result on Integrity


We will make use of the following known result about integrity:

Theorem

Let G be a k -regular Graph of order n . Then

$$k + 1 \leq I(G) \leq n - \text{ind}(G) + 1$$


Example: X_{10}



$$t(X_{10}) = 1$$

$$l(X_{10}) = 6$$

Example: X_{10}



$$t(X_{10}) = 1$$

$$l(X_{10}) = 6$$

Before going on with the results on integrity we will write the bounds for integrity in the language of Unitary Cayley Graphs:

$$\varphi(n) + 1 \leq I(X_n) \leq n - \frac{n}{p_1(n)} + 1$$

Theorem

Let p be a prime number and $k \in \mathbb{Z}^+$. Then
 $I(X_{p^k}) = p^k - p^{k-1} + 1$

Proof.

We know that

$$I(X_{p^k}) \geq \varphi(p^k) + 1 = p^k - p^{k-1} + 1$$

But we also have that:

$$\begin{aligned} I(X_{p^k}) &\leq p^k - \frac{p^k}{p} + 1 \\ &= p^k - p^{k-1} + 1 \end{aligned}$$

Thus $I(X_{p^k}) = p^k - p^{k-1} + 1$ as we wanted to prove. □

Theorem

Let p be a prime number and $k \in \mathbb{Z}^+$. Then
 $I(X_{p^k}) = p^k - p^{k-1} + 1$

Proof.

We know that

$$I(X_{p^k}) \geq \varphi(p^k) + 1 = p^k - p^{k-1} + 1$$

But we also have that:

$$\begin{aligned} I(X_{p^k}) &\leq p^k - \frac{p^k}{p} + 1 \\ &= p^k - p^{k-1} + 1 \end{aligned}$$

Thus $I(X_{p^k}) = p^k - p^{k-1} + 1$ as we wanted to prove. □

Theorem

Let $p < q$ be prime numbers, then $I(X_{pq}) = pq - q + 1$

Proof.

Let S be a vertex-cut set such that $|S| < pq - q$. We have that $k(X_{pq} - S) = 2$. Using the pigeonhole principle we have that:

$$m(X_{pq} - S) \geq \frac{pq - |S|}{2}$$

And so we have, using that $p < q$ and that, as S is a vertex-cut, $|S| \geq \kappa(X_{pq}) = (p-1)(q-1)$:

$$\begin{aligned} |S| + m(X_{pq} - S) &\geq |S| + \frac{pq - |S|}{2} \\ &\geq \frac{|S| + pq}{2} \geq \frac{(p-1)(q-1) + pq}{2} \end{aligned}$$

Proof.

$$\begin{aligned} |S| + m(X_{pq} - S) &\geq \frac{(p-1)(q-1) + pq}{2} \\ &\geq pq - \frac{p+q}{2} + \frac{1}{2} > pq - q \end{aligned}$$

Assume now that $|S| \geq pq - q$, then:

$$|S| + m(X_{pq} - S) \geq pq - q + 1$$

And so $I(X_{pq}) = pq - q + 1$. □

Notice that in both cases the integrity is equal to the upper bound $n - \frac{n}{p_1(n)} + 1$, we conjecture that this is the case for every Unitary Cayley Graph.

GRACIAS!!!!!!!

Bibliography

-  N. Biggs, *Algebraic graph theory, Second Edition*, Chematical library. Cambridge University Press, 1993.
-  G. Chartrand and L. Lesniak, *Graphs and Digraphs, Third Edition*, Chapman and Hall, 1996.
-  C. Godsil and R. Royle, *Algebraic graph theory*, Graduate Text in Mathematics. Springer, 2001.
-  W. Klotz and T. Sander, *Some properties of unitary Cayley graphs*, The Electronic Journal of Combiantorics 14 (2007), R45, pp. 1-12.